

Multiply subtractive generalized
Kramers-Kronig relations: application on third
harmonic generation susceptibility on
polysilane

Valerio Lucarini

lucarini@mit.edu

Jarkko J. Saarinen and Kai-Erik Peiponen

Department of Physics, University of Joensuu,

P. O. Box 111, FIN-80101 Joensuu, Finland

February 2, 2008

February 2, 2008

Abstract

We present multiply subtractive Kramers-Kronig (MSKK) relations for the moments of arbitrary order harmonic generation sus-

ceptibility. Using experimental data on third-harmonic wave from polysilane, we show that singly subtractive Kramers-Kronig (SSKK) relations provide better accuracy of data inversion than the conventional Kramers-Kronig (K-K) relations. The fundamental reason is that SSKK and MSKK relations have strictly faster asymptotic decreasing integrands than the conventional K-K relations. Therefore SSKK and MSKK relations can provide a reliable optical data inversion procedure based on the use of measured data only.

1 Introduction

In linear optical spectroscopy Kramers-Kronig (K-K) dispersion relations [?] have widely been exploited in data inversion both in absorption and reflection spectroscopy [?]. The characteristic integral structure of K-K relations require the knowledge of the spectrum at a semi-infinite angular frequency range. Unfortunately, in practical spectroscopy only finite spectral range can be measured. Moreover, technical difficulties in gathering information especially in nonlinear optical spectroscopy on a sufficiently wide spectral range make hard even the application of the K-K relation. The extrapolations in K-K analysis, such as the estimation of the data beyond the measured spectral range can be a serious source of errors [?, ?]. Recently, King [?] presented an efficient numerical approach to the evaluation of the K-K relations. Nevertheless, the problem of data fitting is always present in regions outside the measured range.

In the context of linear optical spectroscopy singly [?] (SSKK) and multiply [?] (MSKK) subtractive K-K relations have been utilized in order to avoid the limitations due to finite-range data. Only recently they have been proposed in the context of nonlinear optics and especially for harmonic generation susceptibility [?]. The idea behind the subtractive K-K is that the inversion of the real (imaginary) part of n th order harmonic generation nonlinear susceptibility can be greatly improved if we have one or more anchor points, i.e. a single or multiple measurement of the imaginary (real) part for a set of wavelengths. Recently, we presented generalized K-K relations and showed their validity on polymer [?]. In this paper we present for the first time the theory for multiply subtractive generalized K-K for the moments of n th order harmonic generation susceptibility. In addition, we show using data of polysilane that the success of data inversion of the moments relations of third-order harmonic generation susceptibility is better already in the case of SSKK than for conventional K-K. The results of the latter were presented in Ref. [?]. This SSKK is important in the sense that usually it is rather difficult to obtain reliable information on nonlinear susceptibility at multiple anchor points.

2 Multiply subtractive Kramers-Kronig relations for the moments of the harmonic generation susceptibility

We first present the theory related to moments of n th order harmonic generation susceptibility. Then we deal with a function $\omega^{2\alpha}\chi^{(n)}(n\omega; \omega, \dots, \omega)$, where ω is the angular frequency, $0 \leq \alpha \leq n$, and $\chi^{(n)}(n\omega)$ is the tensor element of the nonlinear susceptibility. From now on we make use of the abbreviation $\chi^{(n)}(n\omega; \omega, \dots, \omega) = \chi^{(n)}(n\omega)$. Next we generalize the result of Palmer *et al.* [?] and Lucarini *et al.* [?] to hold also for the moments of the harmonic generation susceptibility. With the aid of mathematical induction (for a rigorous proof see Appendix A in Ref. [?]) we can derive multiply subtractive K-K relations for the real and imaginary parts of as follows:

$$\begin{aligned}
\omega^{2\alpha}\text{Re}\{\chi^{(n)}(n\omega)\} = & \left[\frac{(\omega^2 - \omega_2^2)(\omega^2 - \omega_3^2) \cdots (\omega^2 - \omega_Q^2)}{(\omega_1^2 - \omega_2^2)(\omega_1^2 - \omega_3^2) \cdots (\omega_1^2 - \omega_Q^2)} \right] \omega_1^{2\alpha}\text{Re}\{\chi^{(n)}(n\omega_1)\} + \cdots \\
& + \left[\frac{(\omega^2 - \omega_1^2) \cdots (\omega'^2 - \omega_{j-1}^2)(\omega'^2 - \omega_{j+1}^2) \cdots (\omega'^2 - \omega_Q^2)}{(\omega_j^2 - \omega_1^2) \cdots (\omega_j^2 - \omega_{j-1}^2)(\omega_j^2 - \omega_{j+1}^2) \cdots (\omega_j^2 - \omega_Q^2)} \right] \omega_j^{2\alpha_j}\text{Re}\{\chi^{(n)}(n\omega_j)\} + \cdots \\
& + \left[\frac{(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2) \cdots (\omega^2 - \omega_{Q-1}^2)}{(\omega_Q^2 - \omega_1^2)(\omega_Q^2 - \omega_2^2) \cdots (\omega_Q^2 - \omega_{Q-1}^2)} \right] \omega_Q^{2\alpha}\text{Re}\{\chi^{(n)}(n\omega_Q)\} \\
& + \frac{2}{\pi} [(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2) \cdots (\omega^2 - \omega_Q^2)] \text{P} \int_0^\infty \frac{\omega'^{2\alpha+1} \text{Im}\{\chi^{(n)}(n\omega')\}}{(\omega'^2 - \omega^2) \cdots (\omega'^2 - \omega_Q^2)} d\omega',
\end{aligned} \tag{1}$$

$$\begin{aligned}
\omega^{2\alpha-1}\text{Im}\{\chi^{(n)}(n\omega)\} &= \left[\frac{(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2) \cdots (\omega^2 - \omega_Q^2)}{(\omega_1^2 - \omega_2^2)(\omega_1^2 - \omega_3^2) \cdots (\omega_1^2 - \omega_Q^2)} \right] \omega_1^{2\alpha-1}\text{Im}\{\chi^{(n)}(n\omega_1)\} + \cdots \\
&+ \left[\frac{(\omega^2 - \omega_1^2) \cdots (\omega^2 - \omega_{j-1}^2)(\omega^2 - \omega_{j+1}^2) \cdots (\omega^2 - \omega_Q^2)}{(\omega_j^2 - \omega_1^2) \cdots (\omega_j^2 - \omega_{j-1}^2)(\omega_j^2 - \omega_{j+1}^2) \cdots (\omega_j^2 - \omega_Q^2)} \right] \omega_j^{2\alpha-1}\text{Im}\{\chi^{(n)}(n\omega_j)\} + \cdots \\
&+ \left[\frac{(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2) \cdots (\omega^2 - \omega_{Q-1}^2)}{(\omega_Q^2 - \omega_1^2)(\omega_Q^2 - \omega_2^2) \cdots (\omega_Q^2 - \omega_{Q-1}^2)} \right] \omega_Q^{2\alpha-1}\text{Im}\{\chi^{(n)}(n\omega_Q)\} \\
&- \frac{2}{\pi} [(\omega^2 - \omega_1^2)(\omega^2 - \omega_2^2) \cdots (\omega^2 - \omega_Q^2)] \text{P} \int_0^\infty \frac{\omega'^{2\alpha}\text{Re}\{\chi^{(n)}(n\omega')\}}{(\omega'^2 - \omega^2) \cdots (\omega'^2 - \omega_Q^2)} d\omega',
\end{aligned} \tag{2}$$

with $0 \leq \alpha \leq n$ and P denoting the principal value integration. Here, ω_j with $j = 1, \dots, Q$ denote the anchor points. Note that the anchor points in Eqs. (1) and (2) need not to be the same. We observe that the integrands in MSKK relations presented in Eqs. (1) & (2) have remarkably stronger asymptotic decrease, as a function of angular frequency, than the conventional K-K relations. The integrands of the conventional K-K relations given for the n th order harmonic generation susceptibility converge proportional to $\omega^{-(2n+2)}$ at high angular frequencies, whereas the integrands in MSKK relations converge in relation to $\omega^{-(2n+2+2Q)}$. Therefore, one can expect that the limitations related to the presence of an experimentally unavoidable finite angular frequency range are thus relaxed, and the precision of the integral inversion is then enhanced. According to Palmer *et al.* [?] the anchor points should be chosen inside the measured spectral range. Furthermore,

their study showed that accurate data inversion is possible when the anchor points are chosen near to the zeros of the Q th order Chebyshev polynomial of the first kind. In linear optical spectroscopy it is usually rather easy to get information of the optical constants such as real refractive index and absorption coefficient of the medium at various anchor points. However, in the field of nonlinear optics it may be problematic to obtain the real and imaginary parts of the nonlinear susceptibility at various anchor points. Therefore, in the present study we wish to emphasize that the information even at a single anchor point reduces the errors caused by finite data range by utilization of SSKK. Now the choice of the location of the anchor point is not so critical. Nevertheless, the Chebyshev zeros accumulate at the ends of the spectral range. Therefore, we have chosen the anchor point near one end of the data range. From Eqs. (1) & (2) we obtain for the moments of the third-order harmonic generation susceptibility the following SSKK relations

$$\omega^{2\alpha}\text{Re}\{\chi^{(3)}(3\omega)\} = \omega_1^{2\alpha}\text{Re}\{\chi^{(3)}(3\omega_1)\} + \frac{2(\omega^2 - \omega_1^2)}{\pi} \text{P} \int_0^\infty \frac{\omega'^{2\alpha+1} \text{Im}\{\chi^{(3)}(3\omega')\}}{(\omega'^2 - \omega^2)(\omega'^2 - \omega_1^2)} d\omega', \quad (3)$$

$$\omega^{2\alpha-1}\text{Im}\{\chi^{(3)}(3\omega)\} = \omega_1^{2\alpha-1}\text{Im}\{\chi^{(3)}(3\omega_1)\} - \frac{2(\omega^2 - \omega_1^2)}{\pi} \text{P} \int_0^\infty \frac{\omega'^{2\alpha} \text{Re}\{\chi^{(3)}(3\omega')\}}{(\omega'^2 - \omega^2)(\omega'^2 - \omega_1^2)} d\omega', \quad (4)$$

where ω_1 is the anchor point.

3 Application of SSKK relations to experimental data of third harmonic generation on polysilane

We consider here the experimental data of Kishida *et al.* [?] related to third harmonic susceptibility of polysilane at energy range 0.4–2.5 eV. In Figs. 1a and 2a are shown the inverted data obtained using Eqs. (3) & (4). In the case of Fig. 1a the anchor point was selected at the lower boundary of the data range. The data shown in Fig. 1a for the real part of the nonlinear susceptibility has an almost perfect match with the measured spectrum for $0 \leq \alpha \leq 2$, while for $\alpha = 3$ the finite range causes disagreement between the measured and retrieved data at low energy range of the spectrum. Nevertheless, the spectrum is well retrieved at high energy range. In the case $\alpha = 4$ the theory does not predict convergence, but we have still reasonable agreement between experimental and inverted curve at the high energy range. If we compare the curves of Fig. 1a with those of Fig. 1b obtained using the conventional K-K relations we observe that in Fig. 1b the curve already for $\alpha = 2$ is dramatically departing from the true one at the low energy range. Thus, the SSKK relations provide a better means than the conventional K-K relations to invert the moments data.

In the case of the imaginary part of the nonlinear susceptibility the anchor point was selected at a location of the low energy side of the main spectral

feature, just out of the resonance. Also in the case of the imaginary part we observe that the SSKK procedure provides more precise data inversion than the conventional K-K relation (for comparison see Fig. 2b). In Fig. 2a the agreement between measured and retrieved data is excellent for $0 \leq \alpha \leq 2$ while for $\alpha = 3$ we have a good agreement except for photon energy ≤ 0.7 eV. In the theoretically non-converging $\alpha = 4$ case, we somewhat surprisingly still have a good performance of the SSKK procedure at high energy range. Heuristically we can interpret this result as a manifest of the faster asymptotic decrease of the integrands realized when SSKK relations are utilized instead of K-K relations.

4 Conclusions

We have given, as far as we know, for the first time multiply subtractive Kramers-Kronig relations for the moments of the arbitrary order harmonic generation susceptibility. According to the calculations, which are based on experimental data of polysilane, we have observed how an independent measurement of the unknown part of the complex third-order (arbitrary-order) harmonic generation susceptibility for a given frequency relaxes the limitations imposed by the finiteness of the measured spectral data. The fundamental reason is that SSKK and MSKK relations have strictly faster asymptotic decreasing integrands than the conventional K-K relations. Thus SSKK and MSKK relations can provide a reliable data inversion procedure

based on the use of measured data only. That is to say no extrapolations of the data are needed beyond the measured range. We have demonstrated that even SSKK relations yield more precise data inversion, using a single anchor point, than conventional K-K relations. Naturally it is possible to exploit MSKK procedure if higher precision is required. However, the measurement of data at multiple anchor points may be experimentally tedious.

Acknowledgments

The authors wish to thank Dr. Hideo Kishida and Prof. Takao Koda for providing the measured data on polysilane, and Prof. Franco Bassani for having suggested us to analyze this problem. Authors J.J.S. and K.-E.P wish to express their gratitude to the Academy of Finland for financial support. One of the authors (JJS) wishes to thank the Nokia foundation for a grant.

Figure captions

Figure 1: Efficacy of SSKK relation (a) versus the conventional K-K relation (b) in retrieving $\text{Re}\{\chi^{(3)}(3\omega; \omega, \omega, \omega)\}$ on polysilane.

Figure 2: Efficacy of SSKK relation (a) versus the conventional K-K relation (b) in retrieving $\text{Im}\{\chi^{(3)}(3\omega; \omega, \omega, \omega)\}$ on polysilane.




Figure 1: Lucarini, Saarinen, and Peiponen




Figure 2: Lucarini, Saarinen, and Peiponen